

The association between stress attributed to information systems and the experience of workplace aggression : a cross-sectional survey study among Finnish physicians

Urnberg, Heidi

2022-05-31

Urnberg , H , Gluschkoff , K , Saukkonen , P , Elovainio , M , Vänskä , J & Heponiemi , T
2022 , ' The association between stress attributed to information systems and the
experience of workplace aggression : a cross-sectional survey study among Finnish
physicians ' , BMC Health Services Research , vol. 22 , no. 1 , 724 . <https://doi.org/10.1186/s12913-022-08116-w>

<http://hdl.handle.net/10138/348149>

<https://doi.org/10.1186/s12913-022-08116-w>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

RESEARCH

Open Access



The association between stress attributed to information systems and the experience of workplace aggression: a cross-sectional survey study among Finnish physicians

Heidi Urnberg^{1*}, Kia Gluschko¹, Petra Saukkonen², Marko Elovainio¹, Jukka Vänskä³ and Tarja Heponiemi¹

Abstract

Background: Physicians commonly suffer from workplace aggression and its negative consequences. Previous studies have shown that stressors such as job demands increase the risk of inappropriate treatment at workplace. Poorly functioning, and constantly changing information systems form a major work stressor for physicians. The current study examined the association between physicians' stress attributed to information systems (SAIS) and their experiences of workplace aggression. Workplace aggression covered physical and non-physical aggression, perpetrated by coworkers, patients, patient's relatives, or supervisors.

Methods: A cross-sectional survey study was conducted. The participants included 2786 physicians (67.4% women) who were sampled randomly from the registry of Finnish Medical Association, which covers almost all of the Finnish physician population. First, bivariate associations were studied among participant characteristics, SAIS and workplace aggression. Logistic regression analysis was then used to further determine how SAIS was associated with the likelihood of experiencing different types of aggression.

Results: Higher levels of SAIS were associated with higher likelihood of aggression with regard to all types of aggression, except non-physical aggression perpetrated by patients or relatives. The demographic factors (work-sector, gender, age) did not have a noticeable influence on the association between SAIS and aggression.

Conclusions: The present results build on previous evidence on the prevalence of SAIS and its negative effects on healthcare workers. Since SAIS may increase the risk of experiencing aggression, it is possible that SAIS also endangers the wellbeing of physicians and thereby the quality of patient care. Resourcing time and training during introduction of a new IS could alleviate time pressure and thus stress attributed to managing new information systems. The role of organizational climate and general workload in arousing SAIS and aggression should be examined in future studies.

Keywords: Aggression, Violence, Information systems, SAIS, Job demand, Work stress, Physicians

Background

Workplace aggression is a common and concerning occurrence within healthcare [1–4]. Healthcare professionals, including physicians, have been found to have a heightened risk of experiencing workplace aggression compared to other service workers [5], with estimated one tenth to two thirds of healthcare workers having

*Correspondence: heidi.urnberg@helsinki.

¹ Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, Finland
Full list of author information is available at the end of the article



© The Author(s) 2022. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

encountered aggression [25–27]. It is generally agreed considering that work stress among healthcare workers is becoming increasingly common [10]. Indeed, aggression [5], but the problem is prevalent among physicians as well [24]. The prevalence of workplace aggression is likely even higher than reported because these incidents tend to be underreported [24], which presents an alarming picture of the state of experiencing workplace aggression in healthcare. Because the prevalence of workplace aggression depends on its conceptualization, it is challenging to compare prevalence estimates across different studies. Healthcare data, e.g. electronic health records, which can be used for storing, sharing, searching and retrieving digital patient information [15]. In theory, information systems can offer improvements in quality of care and access to patient information remotely [15], but it is still debated whether information systems actually hinder physicians' physical assaults, harassment, bullying, intimidation, threatening, and obscene behaviors [6].

Workplace aggression can be physical or non-physical in nature and it can manifest for example as verbal abuse, physical assaults, harassment, bullying, intimidation, threatening, and obscene behaviors [6]. Consequently, there is an abundance of proposed constructs of workplace aggression seeking to define the phenomenon. However, many of these constructs lack in definition, well-being [11,14,17,18]. Physicians' complaints regarding workplace aggression often include aspects related to form attributes. Therefore, the definition of "workplace aggression" in this study will encompass simply all these several IS-related factors have been directly associated with manifestations and behaviors towards employees that can result in psychological, social or physical harm to the victim, as suggested in previous literature [8]. In healthcare environments, violence arises typically from patient interactions, but much of the non-physical violence experienced in the healthcare workplace is perpetrated by colleagues and superiors as well [25–27]. However, studies examining workplace aggression have focused mostly on the aggression carried out by patients.

Workplace aggression has consequences for both the well-being of healthcare professionals and quality of patient care. Experiencing workplace aggression is associated with negative psychological outcomes for the healthcare worker (e.g. burnout, anxiety, depression), exhaustion, a common consequence of stress [23], can lead to depersonalization, and subsequently, to negative behavioral changes toward patients and colleagues [1,3,7]. Instances of workplace aggression have also been associated with worse quality of patient care [24]. Secondly, changing, difficult, and poorly functioning information systems may constitute an especially frustrating and stressful work-related stressor for physicians because it may hinder their work, create additional time-pressure, and interfere with patient interaction. Finally, workplace aggression may also have organizational consequences, as SAIS is likely to arouse frustration and negative affect, which may create aggressive inclinations and aggressive behavior (frustration-aggression theory) [25]. As workplace aggression is associated with absences from work [1,4] and the negative psychological consequences resulting from encountering aggression can cause some physicians to leave the organization altogether [9].

Given that workplace aggression has severe consequences, it is important to understand factors that may increase its risk. Work-related stress has been recognized as one of the factors that might increase the risk of suffering from workplace aggression from both patients and other healthcare staff [5–7]. It is especially concerning that the negative affect of a physician might

arouse negative affect in others. Indeed, it has been found that the mood of a healthcare worker is associated with the mood of patients [28] and other healthcare workers [29, 30].

Because workplace aggression is a major concern for the wellbeing of physicians and quality of patient care, it is critical to understand risk factors that are associated with its occurrence. Previous research has shown that stressful work characteristics increase the risk of workplace aggression, potentially by inducing frustration and negative affect. Information systems have been consistently recognized as a major stressful work characteristic for physicians. However, previous studies have not, to our knowledge, examined the association between SAIS and workplace aggression. Therefore, the current study examined the association between physicians' SAIS and their experiences of workplace aggression (physical and non-physical) perpetrated both by patients and health

care staff. This study aims to answer the following main study question:

Are physicians who experience higher levels of SAIS more likely to encounter workplace aggression?

Based on the main study question, following questions were formed:

- 1 Is SAIS associated with non-physical and physical aggression?
- 2 Is SAIS associated with non-physical aggression perpetrated both by patients and relatives, and co-workers and superiors?

Methods

Participants and procedure

Data for this study were drawn from the cross-sectional Finnish Physicians' Working Conditions and Health 2019 source register of the Finnish Medical Association, which covers almost all of the Finnish physician population. The participants who were asked to participate to current study (n = 8000, 37.8% of the Finnish physician population) were approached via email or letter if email was unavailable. A total of 3513 (response rate 43.9%) physicians completed the questionnaire. The representativeness of the sample was assessed by comparing sample and population characteristics. The differences were so small that it was judged that inferences about the Finnish physician population can be made (see, for example [31]).

Physicians who were actively practicing were included in the study (n = 2944) in order to capture the current state of

SAIS and work aggression. The physicians in the sample practiced medicine either in the public sector (covering health centers and hospitals) or in the private sector, such as in private clinics. Public healthcare is available to all permanent residents in Finland regardless of their financial situation. To enable comparable analyses, those who had incomplete demographic information (age, sex or working sector) or missing study variables were removed, resulting in a final analytic sample of 2786 physicians. In this final sample, 67.4% of the participants were female, with a mean age of 45.4 years (standard deviation 11.2), and the majority worked in the public sector (76.2%).

Measures

The questionnaire items for the study variables are presented in the additional file 1.

Workplace aggression

Experiences of both non-physical and physical workplace aggression were measured.

Non-physical workplace aggression was assessed with the following question: "Non-physical violence is defined by ongoing, repeating bullying, oppression or offensive behavior. Do you experience or have you experienced non-physical violence or bullying in your work during the last 12 months?" If a participant answered positively, the origin or perpetrator of aggression was also inquired and the participant could choose any number of the four provided options: coworkers, patients, patient's relatives, and supervisors. From these answers, three dichotomous variables for non-physical aggression were formed: non-physical aggression from any source (yes; 0 no); non-physical aggression from co-workers or superiors (1 yes; 0 no); and non-physical aggression from patients or their relatives (1 yes; 0 no). There were physicians who reported that they had not experienced non-physical aggression, but still reported a specific source of non-physical aggression. These answers were re-coded as "has experienced non-physical aggression". This measure has previously been associated with reduced job satisfaction [31].

Physical workplace aggression was assessed with the following question: "Have you been exposed or threatened via email or letter if email was unavailable with physical violence during the last 12 months?" Participants could choose from three options: "No" (1), "I've only been threatened" (2), "I've also been exposed to violence" (3). A dichotomous variable for physical aggression was formed (1 yes (options 2 and 3, at least threatened with violence); 0 no). This measure has previously been used and associated with increased turnover intention and reduced job satisfaction [31].

A variable for experiencing any type of aggression was formed based on variables of non-physical and physical

aggression (1 yes (has experienced either non-physical aggression, physical aggression, or both) (n0).

Stress attributed to information systems (SAIS)

SAIS was assessed with the following question: "How often (during the past half-year period) have you been troubled by, worried about, or stressed about" with respect to following items: i) "constantly changing information systems" and ii) "difficult, poorly functioning IT equipment/ software". Participants were asked to estimate the level of frequency with a Likert scale from 1 (Very often or constantly) to 5 (Very rarely or never). SAIS was measured with the mean of the two items and the scale's reliability was good in the present sample (Spearman-Brown reliability estimate 0.76).

SAIS is a measure of SAIS that has been previously used in a longitudinal study where it showed good reliability [19], and it has also previously been associated with lower work ability, higher levels of distress and lower self-estimated health [32].

Covariates

All analyses were adjusted for gender (male, 2 female), age, and work sector (public, 2 private), because demographic factors have been associated with risk of aggression in some previous studies [2–6, 32].

Statistical analyses

First, the bivariate associations were studied with Pearson correlation for SAIS and age, and with t-tests for SAIS and other demographic variables (gender and work sector). Then, the differences in SAIS, age, gender, and work sector were analyzed with t-tests and -tests in relation to experiencing a type of aggression (any type of aggression, physical, non-physical, non-physical by patients or relatives, non-physical by co-workers or superiors).

Then, logistic regression analysis was used to determine how different levels of SAIS were associated with

the likelihood of experiencing different types of aggression. The independent variable was the mean score for SAIS. Dependent variables included any type of aggression, physical aggression, non-physical aggression, non-physical aggression perpetrated by patients or relatives, non-physical aggression perpetrated by co-workers or superiors. Each of the aggression types was examined separately in their own models. The models were adjusted for demographic variables (gender, age, and work sector). The results for logistic regression were reported with Odds ratios (OR). The analyses were conducted using R version 3.6.3.

Descriptive and bivariate associations

A fifth (23.4%, n = 651) of the physicians had experienced some type of workplace aggression within the last 12 months. Experiencing non-physical aggression (14.0%, n = 389) was somewhat more common than physical aggression (13.0%, n = 362), with 3.6% having experienced both. Experiencing non-physical aggression perpetrated by co-workers and superiors (9.8%, n = 272) was more common than non-physical aggression perpetrated by patients or relatives (5.0%, n = 139). A total of 1.1% (n = 32) physicians reported experiencing aggression both from patients and relatives and co-workers and superiors.

Higher SAIS was significantly associated with older age ($r = 0.08$, $p < 0.001$). Moreover, higher levels of SAIS were reported by females (mean = 3.40, difference = -0.12, $p = 0.003$), and by those working in the public sector (mean = 3.43) compared to those working in the private sector (mean = 3.08, $t = 7.61$, $p < 0.001$).

The differences between those who reported experiencing some type of aggression and those who did not in SAIS and demographic variables are shown in Table 1. Those who reported experiencing any type of aggression reported significantly higher levels of SAIS. Those who reported experiencing any type of aggression or physical aggression were younger. Instead, those that

reported experiencing non-physical aggression or non-physical aggression from co-workers or superiors were older. Those that experienced aggression were more commonly females in all types of aggression except in physical aggression and non-physical aggression perpetrated by patients or relatives. Similarly, those that experienced physical aggression worked more commonly in the public work sector in all types of aggression except non-physical aggression perpetrated by patients and relatives.

The results of logistic regression

According to the results of logistic regression, SAIS was significantly associated with increased probability for encountering aggression with regard to all the aggression types except non-physical aggression perpetrated by patients or relatives (Table 2). SAIS had the strongest association with non-physical aggression, especially when aggression was perpetrated by co-workers or superiors. Overall, the results of unadjusted and adjusted models were quite similar. Model predicted probabilities for experiencing a specific type of aggression at different levels of SAIS are presented in Fig. 1.

Any type of aggression, physical or non-physical, 1 = yes, 0 = no
Physical aggression 0 = 1 at least threatened
Non-physical aggression 0 = 1 yes
Stress attributed to information systems (1–5) low, 5 high
Gender 0 = male, 1 = female
Work sector 1 = public, 2 = private
Bolded p-values indicate statistical significance

Aggression ^a	Physical aggression ^b			Non-physical aggression ^c			Non-physical aggression by patients and relatives ^d			Non-physical aggression by co-workers and superiors ^e										
	No	Yes		No	Yes		No	Yes		No	Yes									
Variable	Mean (sd)	t	p	Mean (sd)	t	p	Mean (sd)	t	p	Mean (sd)	t	p								
SAIS ^g	3.3 (1.04)	3.5 (1.05)	-4.33	<.001	3.33 (1.04)	3.48 (1.06)	-2.51	.01	-4.55	<.001	3.31 (1.04)	3.52 (1.07)	-1.95	.05	3.32 (45.1)	-4.11	<.001			
Age	45.7 (11.3)	44.3 (11.1)	2.7	.007	45.9 (11.2)	41.8 (11.2)	6.5	<.001	45.2 (11.4)	46.8 (10.4)	-2.84	.005	45.4 (11.2)	44.2 (11.2)	1.26	.21	45.1 (11.4)	48.1 (9.81)	-4.71	<.001
Variable	%	t	p	%	t	p	%	t	p	%	t	p	%	t	p	%	t	p		
Gender ^f (women)	65.7	73.3	12.8	<.001	67.2	68.8	0.27	.60	65.8	77.4	19.8	<.001	67.1	74.1	2.64	.10	66.1	80.1	21.52	<.001
Work sector ^h (public)	75.2	82.3	35.44	<.001	74.2	89.2	38.27	<.001	75.1	82.5	9.65	.002	75.9	80.6	1.32	.25	74.5	82.7	6.74	.01


Table 2 The results of logistic regression models

Predictors	Aggression		Physical aggression		Non-physical aggression		Non-physical by patients or relatives		Non-physical by co-workers or superiors	
	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p
Unadjusted										
SAIS	1.22 (1.11 – 1.33)	< .001	1.15 (1.03 – 1.29)	.012	1.29 (1.15 – 1.44)	< .001	1.19 (1.00 – 1.41)	0.05	1.31 (1.15 – 1.49)	< .001
Adjusted										
SAIS	1.18 (1.08 – 1.30)	< .001	1.15 (1.03 – 1.30)	.015	1.23 (1.10 – 1.37)	< .001	1.18 (0.99 – 1.42)	0.06	1.22 (1.07 – 1.40)	.005

Bolded p-values indicate statistical significance

Adjusted: adjusted for gender, age and work sector

n = 2786, Statistical significance of regression coefficients was tested with Wald's test



Discussion

This is a cross-sectional study among Finnish physicians pronounced with non-physical aggression. It showed that higher levels of SAIS were associated with a higher likelihood of being subjected to all types of negative effects of SAIS in healthcare work environment, except non-physical aggression perpetrated by patients or relatives. The association was most pronounced for non-physical aggression by co-workers or superiors. Our results build on the previous knowledge on the role of SAIS in increasing the likelihood of experiencing aggression, and highlight the possible role of SAIS in increasing the

risk of encountering aggression. Poorly functioning and of non-physical aggression. Moreover, acts of physical constantly changing information systems have previously aggression are perceived less morally acceptable than acts been shown to act as stressors for physicians [14,17], of non-physical aggression, which might make non-phys [18], and work stress has been linked with increased risk of aggression a more likely response to SAIS overall. It of workplace aggression [5–7]. However, to our knowledge should be noted that overall, the observed associations edge, this is the first study to inspect the association were weak, which reflects the fact that reasons underlying between SAIS and experiencing workplace aggression instances of aggression are complex and encompass Potential mechanisms behind this association between numerous other factors that could not be examined in SAIS and aggression could, for example, be that SAIS

The current study. arouses frustration, negative affect, and aggressive inclinations in physicians; and the negative affect and inclination perpetrated by patients or relatives, which might be due to contagion and learning. Previous theories of aggression experiencing non-physical aggression by patients or relatives [25–27] support these suggestions. However, since these mechanisms could not be examined in the current study, they should be examined in future studies, preferably in a longitudinal setting.

Because the current study examined cross-sectional data, it is also possible that experiencing aggression increases SAIS, and not the other way. As shown previously, experiencing aggression is associated with negative psychological outcomes (e.g. burnout, emotional exhaustion and depression) for the healthcare professional, and these outcomes have a negative impact on the work functioning and performance of healthcare professionals [36] and overall psychological health [37]. Moreover, problems in work functioning and performance may lead to more mistakes or increased time pressure. In addition, emotionally exhausted physicians are less able to engage in positive teamwork [33]. In turn, teamwork, the workforce, physician workload, and time pressure have been previously associated with higher levels of SAIS [12].

These results accentuate the need to investigate SAIS and workplace aggression more thoroughly on organizational or department level. The demographic factors did not have a noticeable influence on the association between SAIS and aggression. However, work-sector, gender, and age were significantly associated with aggression. Experiencing aggression was more common among females and in the public work sector, which is in line with previous literature (e.g. [3]). However, findings related to age were here somewhat contradictory: experiencing some types of aggression was more common among younger physicians and others among older physicians. Previous findings on aggression and age have similarly been contradictory [2, 6, 38].

The association between SAIS and aggression appeared to be most pronounced with non-physical aggression.

This finding is consistent with the results of a previous study is subject to some limitations, which should be considered when interpreting the results. First, it should be noted that causal direction between SAIS and aggression could not be determined as the current study used cross-sectional data. Secondly, there were limitations related to the study variables. SAIS was measured as a mean of only two items that asked about negative reactions to information systems. It should be noted that our measurement of stress attributed to information systems (defined as “troubled by, worried about or

stressed about") more accurately depicts negative-psyit may potentially be very consequential in the long run psychological reactions or psychological distress attributed[44]. e present results align with previous literature to information systems. us, our definition of stress on the potential negative effects of SAIS. is notion has follows the model set by Posner and colleagues [39] to decisive implications for the future development of information systems, especially since over a third of physicians is characterized by high arousal. As such, it differs from in our study reported suffering from SAIS frequently or definitions used in work stress literature that use work more often, in line with previous literature [14,17, 18]. stress to refer to the process of work-related stimuli lead Improvements to both the wellbeing of physicians and ing not only to negative psychological but behavioral andthe quality of patient care could be gained in the future physical consequences as well [40]. In addition, all they including proper usability design and active participation variables were assessed only on an individual level and of physicians in the development of IS. Healthcare no information was collected of participants' department organizations should offer easy-to-use systems without or organizational belonging. is means that we measure technical problems, as that would promote the work owed subjective, individual-level perceptions of information and release time for patient work. Moreover, resourcing tion systems and workplace aggression, and could notadditional time and training for physicians during introduction examine, for example, whether physicians working induction of a new IS could also alleviate time pressure and a same department or organization had similar perception thus stress attributed to managing new IS. Special care tions. In addition, using self-report measures might have should be taken in ensuring that IS functions more as a introduced common-methods variance in the results. resource than a demand for the healthcare workers in the Since workplace aggression was measured with subjective future. Similar measures should be heavily considered tive reports rather than objective measures, it is also possible that stress due to high workload might influence a reported to cause stress for physicians in other countries physician to perceive the behavior of others as aggressive as well (e.g [17,18].) along with high prevalence rates of (mood-congruity) [41]. Similarly, high workload might workplace aggression in healthcare (e.g [3].). However, increase SAIS by making it impossible to participate inthe current findings warrant more research, especially IS user training, which could result in low IS competence regarding the possible mediating factors between SAIS and frustration. It is also possible that high workload and aggression. e role of organizational climate and induces stress reactions that lower the tolerance towardsgeneral workload in arousing stress and consequent SAIS poorly functioning IS. Future studies with more objective and aggression should be examined as well. Collecting tive measurements of workplace aggression, workloadongitudinal information on the effect of SAIS on different and IS quality could help shed light on these associationsent types of aggression on organizational or department and help disentangle questions related to cause-effectlevel with different professional groups could provide relationships. Lastly, the current study was conductedfurther insights in future research. with physicians, so the results should be applied to other healthcare professionals with caution.

Conclusion

To conclude, the current study provides evidence that higher levels of SAIS are associated with higher likelihood of experiencing aggression at workplace. Because SAIS may increase the risk of experiencing aggression, it is possible that SAIS also endangers the wellbeing of physicians and thereby the quality of patient care. Poorly functioning information systems are, of course, not the only frustrating factors in the healthcare work environment. It has also been shown that, for example, unfair managerial procedures, high job strain and job demands can provoke aggressive behavior [42]. In addition, the effect sizes of our bivariate differences and Odds ratios are small, thus our results should be interpreted with caution. However, even small effects should be considered, since it has been suggested that even though the size of the effect in psychological research would be very small,

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12913-022-08116-w>

Additional file 1. Measures used in the study.

Acknowledgements

We would like to express our great gratitude to all the physicians who responded to the survey.

Authors' contributions

Substantial contribution to study conception and design: HU, KG, PS, ME, JV, TH. Data analysis: HU and KG. Drafting of the manuscript: HU, KG, PS, ME, JV, TH. All authors read and approved the final manuscript.

Funding

This study was funded by the Strategic Research Council at the Academy of Finland (project 327145). The funder did not have any role in the design of the study and collection, analysis, and interpretation of the data or in the writing.

Availability of data and materials

The data that support the findings of this study are available from Finnish Medical Association, but restrictions apply to the availability of these data, which were used under license for the current study, and so are not publicly

available. The pseudonymized questionnaire data can however be shared by request from the authors after approval of the study group and Finnish Medical Association.

Declarations

Ethics approval and consent to participate

Ethical approval (THL/1037/6.02.01/2019) was obtained from the ethics committee of the Finnish Institute for Health and Welfare. All methods were performed in accordance with these guidelines. All the participants were provided with written information about the project and gave their written informed consent by choosing to participate actively in the study by answering the questionnaire.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Psychology and Logopedics, Faculty of Medicine, University of Helsinki, Helsinki, Finland. ²Finnish Institute for Health and Welfare, Helsinki, Finland. ³The Finnish Medical Association, Helsinki, Finland.

Received: 3 February 2022 Accepted: 20 May 2022

References

1. Lancôt N, Guay S. The aftermath of workplace violence among health-care workers: A systematic literature review of the consequences. *Aggress Violent Behav.* 2014;19(5):492–501. <https://doi.org/10.1016/j.avb.2014.07.010>.
2. Liu J, Gan Y, Jiang H, et al. Prevalence of workplace violence against healthcare workers: a systematic review and meta-analysis Systematic review. *Occup Env Med.* 2019;76:927–37. <https://doi.org/10.1136/oemed-2019-105849>.
3. Mento C, Silvestri MC, Bruno A, et al. Workplace violence against healthcare professionals: A systematic review. *Aggress Violent Behav.* 2020;51(March): 101381. <https://doi.org/10.1016/j.avb.2020.101381>.
4. Phillips JP. Workplace Violence against Health Care Workers in the United States. *N Engl J Med.* 2016;374(17):1661–9. <https://doi.org/10.1056/NEJMp1501998>.
5. Cooper CL, Swanson N. Workplace Violence in the Health Sector. Geneva: State of the Art. International Labour Office; 2002.
6. Camerino D, Estryn-Behar M, Conway PM, Isabella B, Maria Van Der Heijden J, Hasselhorn H-M. Work-related factors and violence among nursing staff in the European NEXT study: A longitudinal cohort study. *Int J Nurs Stud.* 2008;45(1):35–50. <https://doi.org/10.1016/j.ijnurstu.2007.01.013>.
7. Magnavita N. Workplace Violence and Occupational Stress in Healthcare Workers: A Chicken-and-Egg Situation-Results of a 6-Year Follow-up Study. *J Nurs Scholarsh.* 2014;46(5):366–76. <https://doi.org/10.1111/jnu.12088>.
8. Herscovis SM. “Incivility, social undermining, bullying...oh my!”: A call to reconcile constructs within workplace aggression research. *J Organ Behav.* 2011;32(2):499–519. <https://doi.org/10.1002/job>.
9. West CP, Dyrbye LN, Shanafelt TD. Physician burnout: contributors, consequences and solutions. *J Intern Med.* 2018;283(6):516–29. <https://doi.org/10.1111/joim.12752>.
10. Grover S, Adarsh H, Naskar C, Varadharajan N. Physician burnout: A review. *J Ment Heal Hum Behav.* 2018;23(2):78. https://doi.org/10.4103/jmhbb.jmhbb_47_19.
11. Shanafelt TD, Dyrbye LN, Sinsky C, et al. Relationship Between Clerical Burden and Characteristics of the Electronic Environment With Physician Burnout and Professional Satisfaction. In: *Mayo Clinic Proceedings*. Vol 91; Elsevier; 2016:836–848. doi:<https://doi.org/10.1016/j.mayocp.2016.05.007>.
12. Heponiemi T, Hyppönen H, Kujala S, et al. Predictors of physicians' stress related to information systems: a nine-year follow-up survey study. *BMC Health Serv Res.* 2018;18(1):284. <https://doi.org/10.1186/s12913-018-3094-x>.
13. Heponiemi T, Kujala S, Vainiomäki S, et al. Usability factors associated with physicians' distress and information system-related stress: Cross-sectional survey. *J Med Internet Res.* 2019;21(11): e13466. <https://doi.org/10.2196/13466>.
14. Melnick ER, Dyrbye LN, Sinsky CA, et al. The Association Between Perceived Electronic Health Record Usability and Professional Burnout Among US Physicians. *Mayo Clin Proc.* 2020;95(3):476–87. <https://doi.org/10.1016/j.mayocp.2019.09.024>.
15. Black AD, Car J, Pagliari C, et al. The Impact of eHealth on the Quality and Safety of Health Care: A Systematic Overview. *PLoS med.* 2011;8(1): e1000387. <https://doi.org/10.1371/journal.pmed.1000387>.
16. Poissant L, Pereira J, Tamblyn R, Kawasumi Y. The Impact of Electronic Health Records on Time Efficiency of Physicians and Nurses: A Systematic Review. *J Am Med Informatics Assoc.* 2005;12(5):505. <https://doi.org/10.1197/jamia.M1700>.
17. Tajirian T, Stergiopoulos V, Strudwick G, et al. The Influence of Electronic Health Record Use on Physician Burnout: Cross-Sectional Survey. *J Med Internet Res.* 2020;22(7):e19274. <https://doi.org/10.2196/19274>.
18. Gardner RL, Cooper E, Haskell J, et al. Physician stress and burnout: the impact of health information technology. *J Am Med Informatics Assoc.* 2019;26(2):106–14. <https://doi.org/10.1093/jamia/ocy145>.
19. Heponiemi T, Hyppönen H, Vehko T, et al. Finnish physicians' stress related to information systems keeps increasing: A longitudinal three-wave survey study. *BMC Med Inform Decis Mak.* 2017;17(1):147. <https://doi.org/10.1186/s12911-017-0545-y>.
20. Kroth PJ, Morioka-Douglas N, Veres S, et al. Association of Electronic Health Record Design and Use Factors With Clinician Stress and Burnout. *JAMA Netw open.* 2019;2(8):e199609. <https://doi.org/10.1001/jamanetworkopen.2019.9609>.
21. Vänskä J, Vainiomäki S, Kaipio J, Hyppönen H, Reponen J, Lääveri T. Potilasjärjestelmät lääkärin työvälineenä 2014: käyttäjäkokemuksissa ei merkittäviä muutoksia. *Lääkärilehti.* 2014;49:3351–8.
22. Johnson PR, Indvik J. Rudeness at Work: Impulse over Restraint. *Public Pers Manage.* 2001;30(4):457–65.
23. van den Tooren M, de Jonge J. Managing job stress in nursing: what kind of resources do we need? *J Adv Nurs.* 2008;63(1):75–84. <https://doi.org/10.1111/j.1365-2648.2008.04657.x>.
24. Winstanley S, Whittington R. Work & Stress Anxiety, burnout and coping styles in general hospital staff exposed to workplace aggression: A cyclical model of burnout and vulnerability to aggression Anxiety, burnout and coping styles in general hospital staff exposed to workplace aggression. *Work Stress.* 2010;16(4):302–15. <https://doi.org/10.1080/026783702100058650>.
25. Berkowitz L. Frustration-aggression hypothesis: examination and reformulation. *Psychol Bull.* 1989;106(1):59.
26. Bandura A. Psychological mechanisms of aggression. *Aggress Theor Empir Rev.* 1983;1:1–40.
27. Hatfield E, Cacioppo JT, Rapson RL. Emotional contagion. *Curr Dir Psychol Sci.* 1993;2(3):96–100.
28. Haas JS, Cook EF, Puopolo AL, Burstin HR, Cleary PD, Brennan TA. Is the professional satisfaction of general internists associated with patient satisfaction? *J Gen Intern Med.* 2000;15(2):122–8. <https://doi.org/10.1046/j.1525-1497.2000.02219.x>.
29. Chang H-Y, Teng C-I, Chu T-L, Chang H-T, Hsu W-H. Impact of nurse agreeableness and negative mood of nursing supervisors on intention to help colleagues. *J Adv Nurs.* 2012;68(3):636–46. <https://doi.org/10.1111/j.1365-2648.2011.05776.x>.
30. Petitta L, Jiang L, Härtel CEJ. Emotional contagion and burnout among nurses and doctors: Do joy and anger from different sources of stakeholders matter? *Stress Heal.* 2017;33(4):358–69. <https://doi.org/10.1002/ami.2724>.
31. Heponiemi T, Kouvonen A, Virtanen M, Vänskä J, Elovainio M. The prospective effects of workplace violence on physicians' job satisfaction and turnover intentions: The buffering effect of job control. *BMC Health Serv Res.* 2014;14(1):19. <https://doi.org/10.1186/1472-6963-14-19>.
32. Kuusio H, Heponiemi T, Aalto A-M, Sinervo T, Elovainio M. Differences in Well-being between GPs, Medical Specialists, and Private Physicians: The Role of Psychosocial Factors. *Health Serv Res.* 2012;47(1):68–85. <https://doi.org/10.1111/j.1475-6773.2011.01313.x>.

